



Biomechanics of periarticular knee muscles after ACL surgery in young patients – comparison of autograft reconstruction and suture-tape augmented primary repair

PAULINA ZALEWSKA^{1*}, ADRIAN GÓRALCZYK², TOMASZ GUSZCZYN³,
KRZYSZTOF HERMANOWICZ⁴, SZCZEPAN PISZCZATOWSKI¹

¹ Institute of Biomedical Engineering, Białystok University of Technology, Białystok, Poland.

² Humana Medica Omeda, Departament of Orthopaedics, Białystok, Poland.

³ The Medical University of Białystok Children's Clinical Hospital, Białystok, Poland.

⁴ ORTIM Orthopaedic Clinic, Białystok, Poland.

Purpose: Last years brought a growing interest in suture-tape augmented ACL repair techniques instead of ACL reconstructions with autografts due to their limited success rates. However, there is a lack of experimental studies comparing the influence of both procedures on knee periarticular muscles biomechanics. The aim of the study was to analyze periarticular muscles torque and their activity during movement for young patients after ACL repair (IB) and ACL reconstruction (ACLR), compare them with each other and with a healthy subjects (Controls). **Methods:** Isokinetic torques of knee flexors and extensors and their EMG activity during gait and drop vertical jump (DVJ) were measured in 48 patients (IB – 20, ACLR – 28) at least one year after surgery and in 25 healthy controls. **Results:** The muscle torques in the operated limb were lower in both group of patients comparing to Controls, but IB scored the lowest values (79% of Controls for extensors, $p = 0.005$; 74% for flexors, $p = 0.001$). Decrease of muscle torques in the contralateral limb occurred only in IB group (84% of Controls for both extensors ($p = 0.045$) and flexors). Increased hamstrings activity were noticed for ACLR comparing to IB and Controls. The balance between activity of muscles acting on the medial and lateral side of the knee for ACLR tilts more towards the lateral side. **Conclusions:** Increased hamstrings activity and changes in balance between laterally and medially located muscles indicate on the increased need for muscle support of knee stability in ACLR patients.

Key words: knee, ACL reconstruction, Internal Brace, muscle activity, muscle torque

1. Introduction

Anterior cruciate ligament (ACL) tear is one of the most common knee injuries, especially in young, active population [3]. Currently, the gold standard of ACL injuries treatment is reconstruction with use of an autograft [22]. However, in literature variable results are being reported regarding patient's satisfaction and revision rates [23]. As a result, there is a growing interest

in ACL primary repair. One of the most promising technique is a suture-tape augmented repair, for example the Internal Brace™ technique (Arthrex GmBH), (IB) [37].

The IB is a repair technique that allows to reattach and reinforce the native ligament, connecting together the proximal and distal parts of torn ACL using a lasso suture and bracing it with a high-strength suture tape [13], [17]. On the contrary, traditional ACL reconstruction leads to removing injured ligament and replacing

* Corresponding author: Paulina Zalewska, Institute of Biomedical Engineering, Białystok University of Technology, Białystok, Poland. E-mail: paulina.zalewska@sd.pb.edu.pl

Received: December 12th, 2024

Accepted for publication: January 30th, 2025

it by autograft. The necessity of harvesting the graft can cause significant changes in the muscle–tendon unit morphology at the donor-site. In this context, IB technique is presented as less invasive [26] and promising method with potential advantages over reconstruction [13]. Theoretically, the muscle–tendon unit remains unaffected as the autograft is not harvested, which makes it possible to achieve less severe consequences regarding periarticular muscle strength. Moreover, IB may possibly accelerate the rehabilitation protocols and allow for earlier return to full activity due to primary stabilization effect of the suture tape [21].

On the other side, periarticular muscles play significant role as the secondary knee stabilizers [1]. As the ACL surgery result in some changes in the knee structure due to unnatural properties of the reconstructed or repaired ligament (differences in its stiffness, length, proprioception, etc.) it is highly probable that this procedure can influence on periarticular muscle functioning and their activity during movement. Theoretically, this effect should be less prominent in the case of ACL repair as the native ligament is maintained, but all these potential advantages of IB method are still not proven experimentally. To the authors' knowledge, there are practically no studies that assess muscles strength and their activity in a comprehensive way after ACL repair. There is only one study, which is focused on gait analysis [4].

Therefore, the aim of presented study was to:

- analyze the periarticular knee muscles torques and their activity during movement in group of young patients after ACL repair with the Internal Brace™ technique (IB) and after ACL reconstruction with hamstring autograft (ACLR), compare them with each other as well as with a healthy cohort,
- answer to the question if IB method has really advantage over ACLR regarding periarticular muscle functioning.

To our knowledge, this is the first study, which addresses the issue, especially concerning group of young IB patients.

2. Materials and methods

2.1. Concept of the research

In this study, two main aspects were analyzed: (1) knee flexors and extensors strength, (2) muscles activity during movement. Muscles strength were evaluated in the isokinetic condition, while muscle activity was recorded during two functional tests: gait and drop vertical jump (DVJ). All tests were performed during one measurement session lasting no longer than 120 minutes to avoid excessive fatigue of the participant. Taking into account suggestions that the intact limb of the patient after surgery cannot always be treated as a control [14], [24], a group of healthy subjects were included in the research.

2.2. Subjects

Twenty-eight individuals after ACL reconstruction with the hamstrings autograft (ACLR), twenty patients after ACL primary repair with suture tape augmentation according to original Internal Brace™ technique (IB), and twenty five healthy subjects (Controls) participated in this study (Table 1). Patients had completed the rehabilitation process and were at least 12 months postoperatively. Participants were matched to groups so that they did not differ significantly regarding age, body mass and height.

Exclusion criteria included concomitant high grade injury of medial collateral ligament (MCL), lateral collateral ligament (LCL) or posterior cruciate ligament (PCL), meniscus repair in the injured knee, history of ligamentous/meniscal injury, or reconstruction on the contralateral knee. Additionally, exclusion criteria included also history of any injuries in contralateral knee, history of any surgeries involving lower limbs as well as any rheumatological or neurological disorders. All patients were treated in the same way, which was de-

Table 1. Descriptive characteristics of participants (F – female, M – male).
Data are presented as mean and standard deviation [mean (SD)]

Group	Controls (<i>n</i> = 25) 11F / 14M	IB (<i>n</i> = 20) 8F / 12M	ACLR (<i>n</i> = 28) 10F / 18M	<i>p</i> -value
Parameter				
Age [years]	23.7 (1.7)	25.0 (10.5)	21.8 (4.8)	–
Height [m]	1.78 (0.09)	1.76 (0.11)	1.74 (0.12)	–
Mass [kg]	72.9 (13.0)	84.0 (14.5)	76.8 (16.2)	–
Time post-surgery [months]	–	28 (15)	30 (18)	–

scribed more precisely in our earlier paper [10]. The most important reason that IB was performed only for those patients with proximal ACL avulsion within 3 weeks after an injury as well as ACL retained good tissue quality (confirmed intraoperatively). Controls were included on the basis of clinical assessment performed by orthopedist, after meeting following criteria: good general health on the day of the study, no major limb misalignment and no injury in lower limbs in the last 12 months. Participants from all groups do sports only recreationally. A written informed consent was obtained from each subject prior to data collection. The protocol of the study was approved by the Ethical Review Board R-I-002/356/2017.

2.3. Instrumentation and data collection

Isokinetic test

Concentric peak torques of knee extensors and flexors were measured in isokinetic conditions using the BIODEX System 4 Pro (Biodex Medical Systems, New York, USA) at angular velocity 120°/s [15]. During testing, the patient was seated in the upright position, while pelvis, chest and thigh of examined limb were stabilized with straps. The starting position was 90° of knee and 85° of hip flexion. Five practice and four recorded repetitions for both knee flexion and extension were performed, recording muscle torque for the operated (IB-op/ACLR-op), non-operated (IB-nop/ACLR-nop) limb and for both limbs for Controls.

Motion capture

For gait analysis, each participant walked barefoot along a 9-metre walkway at a preferred speed. For DVJ each subject stood on 35-cm high box and were instructed to drop down to the floor and then to immediately performed a vertical jump as fast as possible without stopping. During both functional tests five proper trials (all markers were visible and jump was performed after landing without stopping) were recorded. The motion capture system (Qualisys AB, Gothenburg, Sweden) with set of 26 passive reflective markers and four cluster-tracking markers placed on the lower part of the body was used to determine the basic spatiotemporal parameters of gait and DVJ and time course of knee flexion angle.

EMG

Muscle activity of rectus femoris (RF), vastus medialis (VM), vastus lateralis (VL), semitendinosus

(ST), biceps femoris (BF), and two heads of the gastrocnemius muscle (medial (GM) and lateral (GL)) were recorded using surface electromyography (EMG) (Biometrics, Ltd., Newport, United Kingdom). A set of 7 integrated SX230 electrodes were placed on the tested limb [12]. The frequency of EMG data collection was set at 1000 Hz and band-pass filter was set to 20–450 Hz.

2.4. Data reduction and statistical analysis

The peak extensors and flexors torques values were normalized to a subject's body mass [Nm/kg]. Additionally, torque values obtained for ACLR and IB were compared to value obtained for Controls and expressed as %Controls. To evaluate the symmetry of knee muscles strength between limbs, the Limb Symmetry Index (LSI) was calculated for particular muscle groups by division of the peak torque obtained for the operated limb by its value recorded for the non-operated limb [36]. Symmetrical muscle function for peak torques was defined as achieving $LSI \geq 90\%$, based on the recommendation of the European Board of Sports Rehabilitation [35].

The RMS 150 (Root Mean Square, $t = 150$ ms) was used to obtain meaningful amplitude values from raw EMG data. The activity of analyzed muscles were normalized for each subject separately as a percentage of the arithmetic mean of peak EMG values observed for this participant in all analyzed jump repetitions (%nEMG). Additionally, these activity were presented as time series normalized to the gait cycle for gait (GC) and to jump cycle (JC) for DVJ (period between the first and the last feet contact with the ground). The integrated EMG (iEMG), expressing work done by muscle during GC or JC was determined (%nEMG) [7]. Additionally, a balance indices were calculated, by dividing the temporal normalized activity of laterally situated muscle by concurrent normalized activity of medially located muscle (VL/VM, BF/ST, GL/GM, Σ lateral/ Σ medial), as well as balance between activity of the whole quadriceps and hamstrings (Q/H).

All results were averaged, first for each participant and next, for the whole group (IB-op/ACLR-op). Results for Controls were averaged for both limbs. Additionally, GC was divided into stance and swing phases, while JC for landing and propulsion phases. For statistical analysis, the muscle activities were divided at intervals of 5% of the GC or JC.

The data were checked for normality and homogeneity of variance using histograms and Shapiro–Wilk’s test as well as Levene’s test, respectively. The ANOVA or ANOVA Kruskal–Wallis test with the Tukey’s post-hoc test were used to determine differences between groups and limbs. The significance level was set to $\alpha < 0.05$ and all analyses were performed using STATISTICA 13.1 (TIBCO Software, Inc., USA).

3. Results

3.1. Descriptive characteristics of participants, basic spatiotemporal parameters and knee flexion angle for gait and DVJ

No significant differences were found between groups in descriptive characteristics (Table 1) and basic spa-

tiotemporal gait and DVJ parameters (Table 2), with distinction to knee angle value at the last feet contact with the ground during DVJ ($p = 0.022$).

3.2. Knee extensors and flexors torques

Generally patient’s knee muscles in the operated limb were weaker (Table 3) compared to Controls (79%Controls – IB, 87%Controls – ACLR). IB obtained significantly lower values of extensors ($p = 0.005$) and flexors ($p = 0.001$) torques in both limbs, especially comparing to Controls. Simultaneously, extensors torque in ACLR-nop (2.06 Nm/kg) was significantly higher comparing to ACLR-op (1.78 Nm/kg, $p = 0.030$) and even slightly comparing to Controls (101%Controls). In results, LSI value for extensors was greater for IB (96%), while for ACLR was equal to 88%. For flexors, both groups of patients achieved LSI $\geq 90\%$ (IB – 90%, ACLR – 93%).

Table 2. Basic spatiotemporal parameters of gait and DVJ and knee flexion angle for selected time points. Data are presented as mean and standard deviation [mean (SD)]

Parameter		Group	Controls	IB	ACLR	<i>p</i> -value
GAIT	Spatiotemporal parameters	Stance phase [%]	63.0 (1.5)	62.9 (1.7) ^{op}	62.0 (3.0) ^{op}	–
		Stride length [m]	0.71 (0.05)	0.67 (0.05) ^{op}	0.71 (0.08) ^{op}	–
		Stride time [s]	0.53 (0.04)	0.51 (0.05) ^{op}	0.53 (0.04) ^{op}	–
	Flexion angle	KF 0%GC [°]	2.7 (3.8)	3.5 (5.6) ^{op}	0.6 (5.3) ^{op}	–
		KF 1 [°]	22.5 (4.8)	20.6 (6.2) ^{op}	19.4 (5.5) ^{op}	–
		KF 2 [°]	66.1 (4.9)	66.5 (5.9) ^{op}	63.3 (5.7) ^{op}	–
DVJ	Time parameters	<i>T</i> [s]	0.40 (0.08)	0.41 (0.11) ^{op}	0.37 (0.09) ^{op}	–
		<i>T_L</i> [%]	45.9 (2.2)	44.6 (3.5) ^{op}	46.1 (2.3) ^{op}	–
		<i>T_P</i> [%]	54.1 (2.2)	55.4 (3.5) ^{op}	53.9 (2.3) ^{op}	–
	Flexion angle	0% [°]	26.1 (5.0)	28.7 (6.3) ^{op}	25.9 (6.6) ^{op}	–
		Max [°]	83.8 (11.9)	81.6 (8.8) ^{op}	79.1 (9.9) ^{op}	–
		100% [°]	26.0 (5.4)	23.2 (8.8) ^{op}	20.9 (10.0) ^{op}	0.022 (Control vs. ACLR-op)
	Jump height [% body height]		29.1 (5.1)	26.5 (2.9)	28.1 (4.8)	–

op – operated limb, KF 0%GC – angle value at the beginning of GC, KF 1 – the first knee flexion peak, KF 2 – the second knee flexion peak, *T* – time from the first to the last feet contact with the ground, *T_L* – landing phase time as a percentage of *T*, *T_P* – propulsion phase time as a percentage of *T*, 0% – angle value at the first feet contact with the ground, Max – maximum knee flexion angle, 100% – angle value at the last feet contact with the ground, Jump height was normalized as % of participant’s body height.

Table 3. Peak isokinetic knee flexors and extensors torques and LSI.

Muscle torque was normalized to body weight [Nm/kg].

Data are presented as mean and standard deviation [mean (SD)]

	Extensors			Flexors		
Group	mean (SD) [Nm/kg]	%Controls [%]	LSI [%]	mean (SD) [Nm/kg]	%Controls [%]	LSI [%]
Controls	2.04 (0.36)	—	—	1.41 (0.27)	—	—
IB-op	1.61 (0.43)	79	96 (19)	1.05 (0.26)	74	90 (15)
IB-nop	1.73 (0.55)	84		1.19 (0.35)	84	
ACLR-op	1.78 (0.41)	87	88 (15)	1.22 (0.35)	87	93 (15)
ACLR-nop	2.06 (0.51)	101		1.33 (0.40)	94	
<i>p</i> -value	0.005 (IB-op vs. Controls) 0.045 (IB-nop vs. Controls) 0.040 (IB-nop vs. ACLR-nop) 0.030 (ACLR-op vs. ACLR-nop)		—	0.001 (IB-op vs. Controls)		—

IB – anterior cruciate ligament repair, ACLR – anterior cruciate ligament reconstruction, op – operated limb, nop – non-operated limb, %Controls – percentage of the value obtained by Controls, LSI – Limb Symmetry Index.

3.3. Knee muscles activity

Muscles activities during functional tests were presented in Fig. 1 and Table 4.

Quadriceps

Analyzing RF activity in IB group, it was noted significantly higher activity during gait (0–15% GC) and lower during DVJ (40–65% JC). For ACLR, significant differences were observed only during DVJ (lower for 40–65% JC and higher for 85–100% JC). Regarding VL, statistically significant differences were noted only at landing phase of DVJ, where both, IB (10–35% JC and IEMG value, $p = 0.029$) and ACLR (25–45% JC) reached lower activity than Controls. All patients obtained similar VM activity comparing to Controls during DVJ and higher for gait, but statistically significant increase were noted during swing phase (ACLR – 65–75% GC; IB – 65–80% GC and IEMG value, $p = 0.046$).

Hamstrings

Significantly higher BF activity was noted for ACLR during gait, compared both to Controls (0–80% GC) and IB (0–30% GC). During DVJ, higher BF activity in comparison with Controls was noted, both for ACLR (0–30% JC) and IB (0–15% JC). Significantly higher ST activity, compared to Controls, can be observed at the beginning of gait cycle (10–30% GC) and during propulsion phase (75–100% JC) of DVJ for ACLR and IB (75–90% JC). During gait, significantly higher IEMG values were also noted in the case of ACLR,

both for BF and ST, which means higher hamstrings activity in ACLR patients comparing to others.

Gastrocnemius

Both groups of patients obtained significantly higher GL activity during swing phase of GC compared to Controls (IB – 35–45% GC, ACLR – 30–45% GC). During DVJ, IB, in comparison with Controls, obtained significantly higher GL activity for landing phase (0–10% JC) and its lower activity during propulsion phase (60–75% JC). Regarding GM, ACLR obtained significantly lower activity than others during DVJ (approx. 0–15% JC) and also during gait at the beginning (0–10% GC) and at the end (70–100% GC) of GC.

3.4. Knee muscles balance analysis

Analyzing the knee muscles balance (Fig. 2), it can be observed that generally for ACLR balance tilts more towards the lateral side (higher value of the index) compared to Controls and IB, both for particular muscles pairs and for summary Σ lateral/ Σ medial index. As regards knee extensors, the value of VL/VM index for both groups of patients, remains below the level obtained for Controls, both for gait and DVJ. The exception is the second part of the stance phase of GC, where VL/VM index value for ACLR is higher compared to Controls and even significantly higher comparing to IB (30–50% GC). For knee flexors (BF/ST, GL/GM) greater variability of results occurs, but generally ACLR group reached higher values of




Fig. 1. The waveforms of the mean activity of examined muscles during gait (left) and DVJ (right). Data for IB (green), ACLR (red) and Controls (black), expressed as %nEMG are presented as mean and standard deviation. Phases with statistically significant difference were marked with arrows ($\leftrightarrow$) ($p < 0.05$): $\leftrightarrow$ IB/ACLR (blue), $\leftrightarrow$ IB/Controls (green), $\leftrightarrow$ ACLR/Controls (red)

balance indices than Controls, while IB obtained more similar or lower values of balance indexes compared to Controls.

Analyzing balance between quadriceps and hamstrings (Q/H) it can be noted that all patients obtained

lower values of balance index compared to Controls, both for gait and DVJ, which is mainly the results of higher hamstrings activity. For ACLR difference is even statistically significant at the beginning of GC (15–30% GC).

Table 4. IEMG values of knee periarticular muscles during GC and JC [%nEMG] calculated for IB, ACLR and Controls, presented as mean and standard deviation [mean (SD)]

Pair	Group	Gait			DVJ		
		gait cycle	stance phase	swing phase	jump cycle	landing phase	propulsion phase
		mean (SD)	mean (SD)	mean (SD)	mean (SD)	mean (SD))	mean (SD)
RF	Controls	360 (107)	279 (87)	80 (25)	6971 (291)	2464 (334)	4507 (208)
	IB-op	457 (218)	362 (182)	96 (44)	6688 (606)	2221 (467)	4468 (389)
	ACLR-op	462 (185)	340 (142)	121 (74)	6739 (692)	2308 (621)	4431 (409)
	<i>p</i> -value	–	–	0.040 (ACLR-op vs. Controls)	–	–	–
VL	Controls	519 (147)	444 (131)	75 (28)	6812 (264)	2353 (291)	4459 (227)
	IB-op	477 (204)	401 (180)	76 (38)	6438 (1004)	1951 (564)	4487 (662)
	ACLR-op	529 (196)	447 (167)	82 (50)	6605 (524)	2096 (471)	4513 (222)
	<i>p</i> -value	–	–	–	–	0.029 (IB-op vs. Controls)	–
VM	Controls	393 (156)	333 (106)	77 (40)	6877 (331)	2394 (348)	4484 (261)
	IB-op	561 (360)	430 (251)	131 (118)	6738 (548)	2134 (459)	4604 (226)
	ACLR-op	516 (183)	402 (133)	115 (93)	6906 (605)	2406 (609)	4500 (330)
	<i>p</i> -value	–	–	0.046 (IB-op vs. Controls)	–	–	–
BF	Controls	1236 (345)	854 (252)	382 (125)	5624 (1053)	1883 (540)	3741 (643)
	IB-op	1332 (485)	918 (309)	414 (224)	5853 (1132)	1998 (681)	3855 (795)
	ACLR-op	1921 (849)	1350 (596)	571 (308)	6189 (1245)	2272 (750)	3917 (656)
	<i>p</i> -value	0.014 (ACLR-op vs. IB-op) 0.005 (ACLR-op vs. Controls)	0.011 (ACLR-op vs. IB-op) <0.001 (ACLR-op vs. Controls)	0.013 (ACLR-op vs. Controls)	–	–	–
ST	Controls	1378 (602)	938 (437)	440 (189)	6125 (1196)	2742 (824)	3483 (690)
	IB-op	1556 (515)	1061 (424)	495 (156)	6640 (995)	2765 (609)	3875 (712)
	ACLR-op	1786 (605)	1329 (510)	458 (175)	6398 (1011)	2613 (728)	3785 (634)
	<i>p</i> -value	0.041 (ACLR-op vs. IB-op) 0.030 (ACLR-op vs. Controls)	0.038 (ACLR-op vs. IB-op)	–	–	–	–
GL	Controls	969 (322)	808 (269)	161 (72)	5958 (869)	1938 (462)	4021 (478)
	IB-op	1127 (566)	957 (485)	169 (105)	5719 (1060)	1991 (531)	3727 (657)
	ACLR-op	1117 (437)	985 (386)	132 (59)	6028 (945)	2140 (708)	3887 (592)
	<i>p</i> -value	–	–	–	–	–	–
GM	Controls	1262 (491)	1095 (319)	247 (71)	6062 (881)	2304 (394)	3757 (626)
	IB-op	1250 (289)	1068 (242)	182 (117)	6165 (817)	2370 (601)	3795 (391)
	ACLR-op	1215 (565)	1064 (528)	150 (89)	5844 (1268)	1996 (880)	3848 (566)
	<i>p</i> -value	–	–	0.014 (ACLR-op vs. Controls) 0.041 (IB-op vs. Controls)	–	–	–




Fig. 2. Balance indices for VL/VM, BF/ST and GL/GM as well as summary index $\Sigma_{\text{lateral}}/\Sigma_{\text{medial}}$ and Q/H index during gait (left) and DVJ (right) – data for IB (green), ACLR (red) and Controls (black), presented as mean and standard deviation. Phases with statistically significant difference were marked with arrows ($\leftrightarrow$) ($p < 0.05$): $\leftrightarrow$ IB/ACLR (blue), $\leftrightarrow$ IB/Controls (green), $\leftrightarrow$ ACLR/Controls (red)

4. Discussion

The most important finding of our study is that after ACL surgery, despite completed rehabilitation program, periarticular knee muscle have lower ability to generate torque. Apart from that, changes in their activity during gait and DVJ may be observed.

Interestingly, whereas for ACLR muscle weakness was observed only in the operated limb, both lower extremities were affected for IB. As a result, IB reached LSI value above 90%, which can be positive in the aspect of return to sport [35]. However, the decrease of the ability to generate muscle torques is a negative phenomenon and may indicate the presence of muscle strength deficits also in the uninjured limb [14], [24]. Weakness of the quadriceps [5], [6], [11], [25] and hamstrings [5], [8], [20], [29] muscles for ACLR is a generally well-known. Interestingly, we also found significantly reduced muscle torques in both limbs of IB, compared to Controls and ACLR group. Potentially, patients from IB group could had weaker extensors and flexors muscles even before the surgery, compared to ACLR patients. However, as the main criteria used for patient qualification for IB procedure were ACL good tissue quality and short enough period after injury, no patient selection based on knee muscles forces were performed. It seems, it is interesting topic for further examination. At the same time, some studies shown that the higher force produced by quadriceps, the higher loads acting on the ACL [18], so lower extensors torques could potentially protect ligament healing. For flexors this phenomenon is not entirely clear. Müller et al. [21] also noticed significantly lower extensors torques for patients compared to Controls, but no statistical differences between groups for flexors were observed [21].

As it was previously mentioned, some important changes in the muscle activity during movement were observed. It may be partially explained by the need to compensate deficits in the muscle torques. On the other hand, it may result from affected neuromuscular control. Comparing knee flexors and extensors, increased activity of the first group comparing to the second may be observed in ACLR (Figs. 1, 2). These patients obtained noticeable higher hamstrings activity while quadriceps activity remained on the level similar to Controls. In results, Q/H balance index was significantly lower for ACLR compared to Controls. It is known that the hamstrings muscles show synergistic actions to ACL and prevent excessive anterior tibial translation [16], so reduced hamstrings activation is considered to be a risk factor of ACL injury [33]. On

the other hand, the increased hamstrings activity and its duration is a common compensatory mechanism for the ACL absence or persistent laxity [30]–[32]. It may be observed even after the ACL reconstruction, because the muscle volume is reduced, or the stability of the knee has not been fully restored [19]. It is possible that increased hamstrings activity visible in our results confirm the need for providing additional knee joint stabilization, in the sagittal plane, during gait for ACLR patients. However, if increased flexors activity is accompanied by their co-contraction with extensors it may lead to knee stiffness, which is a known phenomenon for ACLR patients [9].

As it was presented above, primarily for ACLR but partly for IB patients also, increase activity of the whole knee flexors group was observed. However, this phenomenon did not occur evenly for BF and ST muscles. As a result, about 1.5-fold higher value of BF/ST index value can be observed in the second part of GC (40–100% GC) for ACLR compared to other groups, which was mainly due to a greater increase in BF activity. BF/ST index value for IB was similar to Controls during gait, which can be a positive sign. Less visible changes can be observed during DVJ, where for landing phase the BF/ST ratio in IB group remains slightly below the value obtained for ACLR, while for propulsion phase BF/ST index was at a similar level in both groups of patients, remaining at a lower level comparing to Controls (Fig. 2).

Taking knee extensors into account, generally both ACLR and IB patients achieved lower VL/VM balance index values than Controls, which was mainly the results of the lower VL activity during gait and DVJ. However, for ACLR opposite phenomenon was observed in the second part of the stance phase (20–60% GC), when VL/VM index value was higher compared to Controls and IB. This effect remain in coincidence with higher value of BF/ST index in this period, and in result summary index Σ lateral/ Σ medial value remains higher for ACLR compared to IB and Controls (30–100% GC). Peel et al. [27], using a musculoskeletal model, noticed that VL has the greatest effect on ACL loading, so lower activity may be an attempt to reduce its effect on the ACL. It is known that during single support phase of GC, the activity of laterally located muscles is necessary to balance the gravitational forces acting on the medial compartment of the knee. This effect supports the ACL's function. Higher activity of the lateral thigh muscles, noticed for ACLR in our research in the second part of the GC, can be interpreted as the need for providing additional knee joint stabilization and ACL support in the frontal plane during gait. As the dynamic knee varus/valgus deformity presented

no significant differences between ACLR and IB group in our research [10], this enhanced activity of the laterally located thigh muscles seems to be efficient enough.

For gastrocnemius actions, similar activity was obtained in both groups of patients during gait, as well as earlier and longer GL activation was noticed compared to Controls. Rhim et al. [28] stated that gastrocnemius reaction time was significantly faster after ACL reconstruction, what makes it possible to increase joint stability. During gait we observed similar GL/GM balance index value for both IB and ACLR, but higher compared to Controls. In the case of DVJ, increase of GL/GM balance index value can be observed for the whole landing phase for ACLR, while in IB group it remains slightly below the level obtained for Controls in the final part of landing phase. However, it is also worth noting that studies do not agree on gastrocnemius importance [34] – some of them suggest a protective role for the ACL [28], but others attribute an antagonistic role to the ACL [2]. However, the time coincidence of the increased GL/GM balance index value for ACLR patients with similar results for VL/VM and BF/ST pairs of muscles indicates the co-operation of the gastrocnemius muscles in supporting the ACL for providing knee stability.

Presented study has some limitations. Our IB group was slightly older (25.0 ± 10.5) than ACLR (21.8 ± 4.8), as well as heavier than others, but not significantly. Additionally, this study include patients at least 12 months post-surgery, so the findings may not be applicable to the early post-operative rehabilitative stages. Moreover, we have not known yet whether presented changes may persist, aggravate or disappear in a long-time follow-up.

5. Conclusions

Changes in muscle strength and their activity may be observed in patients after ACL surgeries in comparison with healthy controls.

First, patients' knee periarticular muscles were weaker in isokinetic test compared to healthy individuals. The greatest strength deficit was observed for IB (by about 20%), and, surprisingly, the loss of muscle torque was observed also in non-operated limb. As the advantage of such situation is possibility to achieve the acceptable LSI value, when for knee extensors in ACLR it remains below proper level.

Secondly, increased hamstrings activity were noticed for ACLR, comparing both to IB and Controls which can be correlated with enhanced importance of

muscle in providing knee joint stabilization. IB patients takes advantage over ACLR in this aspect.

Thirdly, for ACLR the balance leans more towards the lateral side, especially the second part of the gait cycle (30–100% GC). This phenomenon may indicate the increased need for muscle support in providing knee stability in frontal plane for ACLR patients.

Summing up, the answer to the question if IB method has advantage over ACLR regarding periarticular knee muscle biomechanics is not entirely clear, as both methods have some adverse effects as regards muscle strength deficit and their activity during movement.

Conflict of interest statement

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

Research was performed as a part of project WZ/WM-IIB/5/2023 and financed with use of funds for science from Polish Ministry of Science and Higher Education.

References

- [1] ABULHASAN J.F., GREY M.J., *Anatomy and physiology of knee stability*, J. Funct. Morphol. Kinesiol., 2017, 2 (4).
- [2] ADOUNI M., SHIRAZI-ADL A., MAROUANE H., *Role of gastrocnemius activation in knee joint biomechanics: gastrocnemius acts as an ACL antagonist*, Comput. Methods Biomech. Biomed. Engin., 2016, 19 (4), 376–385.
- [3] BRAM J.T., MAGEE L.C., MEHTA N.N., PATEL N.M., GANLEY T.J., *Anterior Cruciate Ligament Injury Incidence in Adolescent Athletes: A Systematic Review and Meta-analysis*, Am. J. Sports Med., 2021, 49 (7), 1962–1972.
- [4] BÜHL L., MÜLLER S., NÜESCH C., BOYER K.A., CASTO E., MÜNDERMANN A., EGLOFF C., *Ambulatory knee biomechanics and muscle activity 2 years after ACL surgery: InternalBraceTM-augmented ACL repair versus ACL reconstruction versus healthy controls*, BMC Musculoskelet. Disord., 2023, 24 (1), 785.
- [5] CZAMARA A., *Moments of muscular strength of knee joint extensors and flexors during physiotherapeutic procedures following anterior cruciate ligament reconstruction in males*, Acta Bioeng. Biomech., 2008, 10 (3), 37–44.
- [6] DUJARDIN D., FONTANIN N., GEFFRIER A., MOREL N., MENSA C., OHL X., *Muscle recovery after ACL reconstruction with 4-strand semitendinosus graft harvested through either a posterior or anterior incision: A preliminary study*, Orthop. Traumatol. Surg. Res., 2015, 101 (5), 539–542.
- [7] EINARSSON E., THOMSON A., SAS B., HANSEN C.L., GISLASON M., WHITELEY R., *Lower medial hamstring activity after ACL reconstruction during running: A cross-sectional study*, BMJ Open Sport Exerc. Med., 2021, 7 (1), 1–5.

- [8] VON ESSEN C., MCCALLUM S., ERIKSSON K., BARENIUS B., *Minimal graft site morbidity using autogenous semitendinosus graft from the uninjured leg: a randomised controlled trial*, Knee Surgery, Sport Traumatol. Arthrosc., 2022, 30 (5), 1639–1645.
- [9] GARCIA S.A., JOHNSON A.K., BROWN S.R., WASHABAUGH E.P., KRISHNAN C., PALMIERI-SMITH R.M., *Dynamic knee stiffness during walking is increased in individuals with anterior cruciate ligament reconstruction*, J. Biomech., 2023, 146, 111400.
- [10] GÓRALCZYK A., ZALEWSKA P., PISZCZATOWSKI S., HERMANOWICZ K., GUSZCZYN T., *No difference in laxity, proprioception and neuromuscular control after suture-tape augmented ACL repair of acute proximal avulsions versus ACL reconstruction using hamstring autografts in young, active population*, J. Exp. Orthop., 2024, 11 (3), e70025.
- [11] HARDY A., CASABIANCA L., ANDRIEU K., BAVEREL L., NOAILLES T., *Complications following harvesting of patellar tendon or hamstring tendon grafts for anterior cruciate ligament reconstruction: Systematic review of literature*, Orthop. Traumatol. Surg. Res., 2027, 103 (8), S245–S248.
- [12] HERMENS H., FRERIKS B., *The state of the art on sensors and sensor placement procedures for surface electromyography*, Roessingh Research and Development, 1997.
- [13] HEUSDENS C.H.W., *Acl repair: A game changer or will history repeat itself? A critical appraisal*, J. Clin. Med., 2021, 10 (5), 1–12.
- [14] HIEMSTRA L.A., WEBBER S., MACDONALD P.B., KRIELLAARS D.J., *Contralateral limb strength deficits after anterior cruciate ligament reconstruction using a hamstring tendon graft*, Clin. Biomech. (Bristol, Avon), 2007, 22 (5), 543–550.
- [15] HUBER R., VIECELLI C., BIZZINI M., FRIESENBICHLER B., DOHM-ACKER M., ROSENHECK T., HARDER L., MAFFIULETTI N.A., *Knee extensor and flexor strength before and after anterior cruciate ligament reconstruction in a large sample of patients: influence of graft type*, Phys. Sportsmed., 2019, 47 (1), 85–90.
- [16] IMRAN A., O'CONNOR J.J., *Control of knee stability after ACL injury or repair: Interaction between hamstrings contraction and tibial translation*, Clin. Biomech., 1998, 13 (3), 153–162.
- [17] MACKAY G.M., BLYTH M.J.G., ANTHONY I., HOPPER G.P., RIBBANS W.J., *A review of ligament augmentation with the InternalBrace™: the surgical principle is described for the lateral ankle ligament and ACL repair in particular, and a comprehensive review of other surgical applications and techniques is presented*, Surg. Technol. Int., 2015, 26, 239–255.
- [18] MANIAR N., COLE M.H., BRYANT A.L., OPAR D.A., *Muscle Force Contributions to Anterior Cruciate Ligament Loading*, Sport Med., 2022, 52 (8), 1737–1750.
- [19] MANTASHLOO Z., LETAFATKAR A., MORADI M., *Vertical ground reaction force and knee muscle activation asymmetries in patients with ACL reconstruction compared to healthy individuals*, Knee Surgery, Sport Traumatol. Arthrosc., 2020, 28 (6), 2009–2014.
- [20] DU MOULIN W., BOURNE M., DIAMOND L.E., KONRATH J., VERTULLO C., LLOYD D., SAXBY D.J., *Shape differences in the semitendinosus following tendon harvesting for anterior cruciate ligament reconstruction*, J. Orthop. Res., 2022, 41 (1), 44–53.
- [21] MÜLLER S., BÜHL L., NÜESCH C., PAGENSTERT G., MÜNDERMANN A., EGLOFF C., *Favorable Patient-Reported, Clinical, and Functional Outcomes 2 Years After ACL Repair and InternalBrace Augmentation Compared With ACL Reconstruction and Healthy Controls*, Am. J. Sports Med., 2023, 51 (12), 3131–3141.
- [22] MURRAY M.M., *Optimizing outcomes of ACL surgery – Is autograft reconstruction the only reasonable option?*, J. Orthop. Res. Off Publ. Orthop. Res. Soc., 2021, 39 (9), 1843–1850.
- [23] PACHE S., DEL CASTILLO J., MOATSHE G., LAPRADE R.F., *Anterior cruciate ligament reconstruction failure and revision surgery: Current concepts*, J. ISAKOS, 2020, 5 (6), 351–358.
- [24] PAIROT DE FONTENAY B., ARGAUD S., BLACHE Y., MONTEIL K., *Contralateral limb deficit seven months after ACL-reconstruction: an analysis of single-leg hop tests*, Knee, 2015, 22 (4), 309–312.
- [25] PALMIERI-SMITH R.M., LEPLEY L.K., *Quadriceps strength asymmetry after anterior cruciate ligament reconstruction alters knee joint biomechanics and functional performance at time of return to activity*, Am. J. Sports Med., 2015, 43 (7), 1662–1669.
- [26] PANG L., LI P., LI T., LI Y., ZHU J., TANG X., *Arthroscopic Anterior Cruciate Ligament Repair Versus Autograft Anterior Cruciate Ligament Reconstruction: A Meta-Analysis of Comparative Studies*, Front. Surg., 2022, 20, 9, 887522.
- [27] PEEL S.A., SCHROEDER L.E., WEINHANDL J.T., *Lower extremity muscle contributions to ACL loading during a stop-jump task*, J. Biomech., 2021, 121, 110426.
- [28] RHIM H.C., LEE J.H., HAN S.B., SHIN K.H., SUH D.W., JANG K.M., *Role of the triceps surae muscles in patients undergoing anterior cruciate ligament reconstruction: A matched case-control study*, J. Clin. Med., 2020, 9 (10), 1–11.
- [29] SANADA T., IWASO H., FUKAI A., HONDA E., YOSHITOMI H., INAGAWA M., *Anatomic Anterior Cruciate Ligament Reconstruction Using Rectangular Bone–Tendon–Bone Autograft versus Double–Bundle Hamstring Tendon Autograft in Young Female Athletes*, Arthrosc. Sport Med. Rehabil., 2021, 3 (1), e47–e55.
- [30] SHANBEHZADEH S., MOHSENI BANDPEI M.A., EHSANI F., *Knee muscle activity during gait in patients with anterior cruciate ligament injury: a systematic review of electromyographic studies*, Knee Surg. Sports Traumatol. Arthrosc., 2017, 25 (5), 1432–1442.
- [31] SHARIFI M., SHIRAZI-ADL A., *Knee flexion angle and muscle activations control the stability of an anterior cruciate ligament deficient joint in gait*, J. Biomech., 2021, 117.
- [32] SHARIFI M., SHIRAZI-ADL A., MAROUANE H., *Computational stability of human knee joint at early stance in Gait: Effects of muscle coactivity and anterior cruciate ligament deficiency*, J. Biomech., 2017, 63, 110–116.
- [33] SØRENSEN B., AAGAARD P., MALCHOW-MØLLER L., ZEBIS M.K., BENCKE J., *Medio-lateral hamstring muscle activity in unilateral vs. Bilateral strength exercises in female team handball players – a cross-sectional study*, Int. J. Sports Phys. Ther., 2021, 16 (3), 704–714.
- [34] STEWART C., SHORTLAND A.P., *The biomechanics of pathological gait – from muscle to movement*, Acta Bioeng. Biomech., 2010, 12 (3), 3–12.
- [35] THOMÉÉ R., KAPLAN Y., KVIST J., MYKLEBUST G., RISBERG M.A., THEISEN D., TSEPIS E., WERNER S., WONDRASCH B., WITVROUW E., *Muscle strength and hop performance criteria prior to return to sports after ACL reconstruction*, Knee Surg. Sports Traumatol. Arthrosc., 2011, 19 (11), 1798–1805.
- [36] WELLSANDT E., FAILLA M.J., SNYDER-MACKLER L., *Limb Symmetry Indexes Can Overestimate Knee Function After Anterior Cruciate Ligament Injury*, J. Orthop. Sports Phys. Ther., 2017, 47 (5), 334–338.
- [37] WILSON W.T., HOPPER G.P., BANGER M.S., BLYTH M.J.G., RICHES P.E., MACKAY G.M., *Anterior cruciate ligament repair with internal brace augmentation: A systematic review*, Knee, 2022, 35, 192–200.